

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07D 235/18, A01N 43/52, C07D 417/10, A01N 43/78, C07D 405/04, 403/04, 405/14, A01N 43/54, C07D 455/04		A1	(11) International Publication Number: WO 99/11627 (43) International Publication Date: 11 March 1999 (11.03.99)
(21) International Application Number: PCT/US98/18586 (22) International Filing Date: 4 September 1998 (04.09.98) (30) Priority Data: 08/924,558 5 September 1997 (05.09.97) US (71) Applicant: ORTHO-MCNEIL PHARMACEUTICAL, INC. [US/US]; U.S. Route 202, Raritan, NJ 08869 (US). (72) Inventors: OHEMENG, Kwasi, Adomako; 4 Shannon Way, Flemington, NJ 08822 (US). NGUYEN, Van, Nhatton; 106 Canal Park, Easton, PA 18042 (US). (74) Agents: CIAMPORCERO, Audley et al.; Johnson & Johnson, 1 Johnson & Johnson Plaza, New Brunswick, NJ 08933 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.	
(54) Title: 2-SUBSTITUTED PHENYL-BENZIMIDAZOLE ANTIBACTERIAL AGENTS			
(57) Abstract			
<p>The invention relates to benzimidazole antibacterial compounds of formula (I) as described herein, pharmaceutical compositions containing the compounds, and methods for their production and use. These compounds are effective in inhibiting the action of a bacterial histidine protein kinase and are useful as anti-infective agents against a variety of bacterial organisms, including organisms which are resistant to other known antibiotics.</p>		<p style="text-align: right;">(I)</p>	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KR	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

2-SUBSTITUTED PHENYL-BENZIMIDAZOLE ANTIBACTERIAL AGENTS

FIELD OF THE INVENTION

5 The invention relates to benzimidazole antibacterial compounds, pharmaceutical compositions containing the compounds, and methods for their production and use. These compounds are effective in inhibiting the action of a bacterial histidine protein kinase and are useful as anti-infective agents against a variety of bacterial organisms, including organisms which are resistant to other known antibiotics.

10

BACKGROUND OF THE INVENTION

It is well established that prokaryotes regulate the transcription of many of their genes in response to changes in the organisms' environment (J. B. Stock, A. M. Stock, 15 and J. M. Mottonen, *Nature*, **344**, 395-400 (1990)). Such regulation is essential if the organism is to adapt itself to survival in a changing environment, and pathogenic bacteria rely on such regulatory systems to enable them to survive within their host's body (J. F. Miller, J. J. Mekalanos, S. Falkow, *Science*, **243**, 1059 (1989)). Chemical compounds that interfere with the regulatory mechanisms would be expected to be useful anti- 20 infective drugs, as they would prevent bacteria from making necessary adaptive changes in their patterns of gene expression.

Virulence, chemotaxis, toxin production, sporulation, and reproduction are examples of the bacterial processes that are under regulatory control, and which could be 25 inhibited by such compounds. The inhibition of one or more of these processes is expected to lead to reduced virulence, a slowing or halting of bacterial growth and reproduction, and even to bacterial cell death if vital functions are interrupted.

For example, it has been shown that *Salmonella* species express certain proteins, under regulatory control and in response to the presence of epithelial cells, which enable them to adhere to and invade intestinal epithelial cells. Bacteria that are unable to synthesize these proteins are avirulent: they cannot cause infection in mice (B. B. Finlay, 5 F. Heffron, S. Falkow, *Science*, 243, 940-943 (1989)). A similar effect would be expected if the genes coding for these proteins were intact, but remained unexpressed.

To accomplish adaptive responses to the environment, bacteria rely on phosphorelay mechanisms, referred to in the art as a "two-component switches." These 10 switches have the net effect of transmitting information from the environment to the cell nucleus, where the information is responded to by the switching on or off of transcription of relevant genes. The first step of this phosphorelay scheme relies on numerous histidine protein kinase (HPK) enzymes. Most of these HPK enzymes are sensor molecules, and respond to stimulation by specific environmental signals by 15 transferring phosphate from ATP to a histidine residue of the HPK protein. Some HPK enzymes are stimulated by the presence of acceptor proteins (described below), the concentration of which are modulated by environmental signals. In either case, this auto-phosphorylation is followed by transfer of the phosphate to an aspartyl residue of one or more acceptor proteins (the second components of the two-component switch), 20 which are either regulators of gene expression (by binding to control regions on DNA, or to the RNA polymerase complex) or are themselves kinases for other acceptor molecules. These secondary acceptors may again be regulatory proteins, or kinases toward yet another protein. This cascade of phosphate from protein to protein eventually results in the phosphorylation of one or more regulatory proteins, which then control 25 gene expression.


Mammalian cells do not, or at least are not presently known to, utilize HPK-driven phosphorelay systems for gene regulation. Thus, compounds which selectively inhibit either the autophosphorylation of the HPK protein, or the phosphotransfer

step(s), or both, would not be expected to have undesirable effects on the host organism, and are promising candidates for antiinfective drugs. The emergence of drug-resistant pathogenic organisms that are resistant to one or more of the currently available drugs has created a need for novel antibiotics, that act by mechanisms unrelated to those of currently available agents, and inhibitors of HPK would fill this need. The presence of multiple HPK-driven systems (over fifty are currently known) in bacteria gives HPK inhibitors a potential advantage over current antibiotics, in that mutations of a single HPK enzyme are unlikely to confer drug resistance to an organism.

Recently, workers in this field reported a method for detecting bacterial "virulence" genes that are selectively expressed when bacteria infect a host (M. J. Mahan, J. M. Slauch, and J. J. Mekalanos, *Science*, **259**, 686-688 (1993)). The potential use of this information in the design of new antibiotics was mentioned, but actual methods of reducing expression of these genes were not described. A preliminary report from another group of workers disclosed inhibitors of the two-component switch controlling alginate gene activation in *Pseudomonas aeruginosa* in an *in vitro* system (S. Roychoudhury *et al.*, *Proc. Nat. Acad. Sci.*, **90**, 965-969 (1993)), but no anti-bacterial activity of the compounds was reported.

SUMMARY OF THE INVENTION

The invention relates to compounds of the Formula I :




Formula I.

wherein:

R₁ is selected from H, OH, (C₁-C₆)alkoxy, and -O(C₁-C₃)COOH;


R₂, R₃ and R₄ are independently selected from H, OH, (C₁-C₁₀)alkyl (branched or
 5 unbranched), (C₁-C₁₀)alkoxy, -O(C₁-C₃)COOH, -O(C₁-C₆)alkylphenyl, -OCH₂(4-
 NO₂phenyl), CF₃, halo, phenyl, substituted phenyl (wherein the substituents are
 selected from halogen, (C₁-C₆)alkoxy, nitro, NH₂, and (C₁-C₆)alkyl), -(C₁-
 C₆)alkylphenyl, nitro, -O(CH₂)_qN((C₁-C₆)alkyl)₂ wherein q is 1-3, -NR₁₇R₁₈, 2-
 thiazole optionally substituted by a (C₁-C₆) alkyl group, -CH=CH-phenyl, pyridinyl,
 10 piperidinyl, -N-phenyl substituted piperidine,
 -O-piperazine, pyrrolinyl, and a moiety of the formula:



wherein n is an integer from 0-2;

R₁₂, R₁₃, R₁₄, R₁₅ and R₁₆ are independently selected from H, nitro, halo,
 15 CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and amino, amidine, sulfonate,
 -C(O)O(C₁-C₆)alkyl, and carboxyl;

or R₁₃ and R₁₄ or R₁₄ and R₁₅ taken together form a fused 5 membered
 heterocyclic ring of the formula:




20 which may be optionally substituted wherein the substituent is selected from
 amino and methyl;

R₁₇ and R₁₈ are independently selected from H, benzyl, phenyl, (C₁-C₆)alkyl,
 CH₂CN, CH₂CH₂CN, or R₁₇ and R₁₈ taken together with the nitrogen to

which they are attached form a 5-6 membered monocyclic heterocyclic ring

optionally containing 1-2 additional nitrogen atoms;


or R_2 and R_3 , or R_3 and R_4 taken together with the phenyl ring to which they are attached form a moiety selected from those of the formulae:



is CH, O or N and R_{20} is (C_1-C_3) alkyl or OH;

R_5 is selected from H, amino and amidino;

R_6 is selected from nitro and moieties of the formula:



10

and



wherein:

R_9 is hydrogen and R_{10} is selected from H or OH or R_9 and R_{10} taken


together form a 5, 6 or 7 membered monocyclic heterocyclic ring optionally

15

containing 1 or 2 additional nitrogens and optionally substituted wherein the

substituent is dimethyl, or R_9 and R_{10} taken together form a bicyclic

heterocyclic ring of the formula;



R_7 is selected from H, amino and nitro;

20

R_8 is selected from H and methyl;

and the pharmaceutically acceptable salts, esters and prodrug forms thereof.

The compounds of the present invention inhibit the autophosphorylation of bacterial histidine kinases; they also inhibit the transfer of phosphate from phosphorylated histidine kinases to the aspartyl residues of the phosphate acceptor proteins involved in regulation of bacterial gene expression. The compounds of the present invention have been found to inhibit the growth of bacteria by the standard method, measurement of minimum inhibitory concentrations (MIC values). The compounds are useful as bacteriostatic and bactericidal agents.

10

DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term "alkyl" means branched or unbranched hydrocarbon chains, such as, for example, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *tert*-butyl, and the like. The alkyl group may be optionally substituted with one or more functional groups commonly attached to such chains, for example, hydroxy, halo, mercapto, thio, cyano, alkylthio, nitro, amino, alkoxy, amido, carboxyl, alkoxycarbonyl, amidino, guanidino and the like.

The term "alkoxy" means the group -OR, where R is an alkyl group as defined above, such as methoxy, ethoxy, *n*-propoxy, *iso*-propoxy, *n*-butoxy, *sec*-butoxy, *iso*-butoxy, *tert*-butoxy and the like.

The term "amino" includes primary, secondary and tertiary amines of the formula -NRR' where R and R' are independently hydrogen or alkyl as defined above.


25

The term "amido" means a group of the formula -C(O)NRR', where R and R' may independently be hydrogen or alkyl as defined above.


The term "prodrug" means an agent which is converted into the parent drug in vivo. Prodrugs may be used to deliver the drug in a manner that is easier or more efficiently delivered. For instance, a drug which is not orally bioavailable may be orally administered in a prodrug form which is orally available, or the prodrug form may be
 5 more soluble for administration intravenously.

Within the group of compounds defined by Formula I, certain subgroups of compounds are preferred. Broadly preferred are those compounds of Formula I wherein:


- 10 R_1 is selected from H, OH, (C₁-C₆)alkoxy, and -O(C₁-C₃)COOH;
 R_2 is selected from H, (C₁-C₁₀)alkyl (branched or unbranched), (C₁-C₁₀)alkoxy, -O(C₁-C₁₀)alkylphenyl, CF₃, halo, phenyl, nitro, pyridyl, and a moiety of the formula:




- R_3 is selected from H, OH, (C₁-C₁₀)alkyl (branched or unbranched), (C₁-
 15 C₁₀)alkoxy,
 -O(C₁-C₆)alkylphenyl, -OCH₂(4-NO₂phenyl), CF₃, halo, phenyl, substituted phenyl (wherein the substituents are selected from halogen, (C₁-C₆)alkoxy, nitro, NH₂, and (C₁-C₆)alkyl), -(C₁-C₆)alkylphenyl, nitro, -O(CH₂)_qN((C₁-C₆)alkyl)₂ wherein q is 1-3, -NR₁₇R₁₈, 2-thiazole optionally substituted by a (C₁-C₆) alkyl group, -CH=CH-
 20 phenyl, -N-phenyl substituted piperidine, -O-piperazine, pyrrolinyl, and a moiety of the formula:



- wherein R₁₇ and R₁₈ are independently selected from H, benzyl, phenyl, (C₁-C₆)alkyl, CH₂CN, CH₂CH₂CN, or R₁₇ and R₁₈ taken together with the nitrogen to which they are attached form a 5-6 membered monocyclic heterocyclic ring optionally containing 1-2 additional nitrogen atoms;
- 5 R₄ is selected from H, (C₁-C₁₀)alkyl (branched or unbranched), (C₁-C₁₀)alkoxy; -O(C₁-C₆)alkylphenyl, CF₃, halo, phenyl, nitro and a moiety of the formula:




or R₂ and R₃ or R₃ and R₄ taken together with the phenyl ring to which they are attached form a moiety selected from those of the formulae:



is CH, O or N and R₂₀ is (C₁-C₃)alkyl or OH;

R₅ is selected from H, amino and amidino;

R₆ is selected from nitro and moieties of the formula:



15


and



wherein:

- R₉ is hydrogen and R₁₀ is selected from H or OH or R₉ and R₁₀ taken together form a 5, 6 or 7 membered monocyclic heterocyclic ring optionally containing 1 or 2 additional nitrogens and optionally substituted wherein the
- 20

substituent is dimethyl, or R₉ and R₁₀ taken together form a bicyclic heterocyclic ring of the formula;



R₇ is selected from H, amino and nitro;

5 R₈ is selected from H and methyl;

R₁₂ is selected from H and nitro;

R₁₃ is selected from H, nitro, halo, CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and amino;


R₁₄ is selected from H, nitro, halo, CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, amino,

amidine, sulfonate, -C(O)O(C₁-C₆)alkyl, and carboxyl;

10 R₁₅ is selected from H, nitro, halo, CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and amino;

R₁₆ is selected from H and nitro;

or R₁₃ and R₁₄ or R₁₄ and R₁₅ taken together form a fused 5 membered heterocyclic ring of the formula:



15 which may be optionally substituted wherein the substituent is selected from amino and methyl;


n is an integer from 0-2;

and the pharmaceutically acceptable salts, esters and prodrug forms thereof.

20 More particularly preferred are the compounds wherein R₁ -R₄ are as described above;


R₅ is H;

R₆ is a moiety of the formula:



wherein:

R₉ is hydrogen and R₁₀ is selected from H or OH or R₉ and R₁₀ taken together form a 5, 6 or 7 membered monocyclic heterocyclic ring optionally containing 1 or 2 additional nitrogens and optionally substituted wherein the
 5 substituent is dimethyl, or R₉ and R₁₀ taken together form a bicyclic heterocyclic ring of the formula;



R₇ and R₈ are hydrogen and R₁₂ to R₁₆ are as described above;
 and the pharmaceutically acceptable salts, esters and prodrug forms thereof.


10

Most preferred are the 5-amidino compounds of Formula I wherein:

R₁ is selected from H, OH, (C₁-C₆)alkoxy;

R₂ is selected from H, (C₁-C₁₀)alkyl (branched or unbranched), (C₁-C₆)alkoxy; -

O(C₁-C₁₀)alkylphenyl, CF₃, halo, phenyl, nitro and a moiety of the formula:




15

R₃ is selected from H, OH, (C₁-C₁₀)alkyl (branched or unbranched), (C₁-C₁₀)alkoxy,

-O(C₁-C₆)alkylphenyl, CF₃, halo, phenyl, substituted phenyl (wherein the substituents are selected from halogen, (C₁-C₆)alkoxy, nitro, NH₂, and (C₁-C₆)alkyl),

20 -(C₁-C₆)alkylphenyl and and a moiety of the formula:



R₄ is selected from H, (C₁-C₁₀)alkyl (branched or unbranched), and -OCH₂phenyl;

R₅ is H;

R₆ is amidine;

5 R₇ is H;

R₈ is H;

n is 0;

R₁₂ is H;

R₁₃ is selected from H, nitro, halo, CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and amino;

10 R₁₄ is selected from H, nitro, halo, CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy and amino;

R₁₅ is selected from H and (C₁-C₆) alkoxy, and amino;


R₁₆ is H;

and the pharmaceutically acceptable salts, esters and prodrug forms thereof.

15 The compounds of the present invention may generally be prepared in accordance with the following schemes, but other synthetic pathways are possible and are within the scope of this invention.

The benzimidazole compounds of the present invention may be prepared in
 20 accordance with Scheme I by one of the following two routes: A) Cyclization of the appropriately substituted diaminophenyl compound with an appropriately substituted aldehyde [(as the bisulphite adduct) in accordance with the procedures disclosed by H. F. Ridley, R. G. W. Spickett and G. M. Timmis; J. Heterocycl Chem 2, 455, 1965] or
 B) Cyclization of the diaminophenyl compound with the appropriately substituted
 25 imidate to give the benzimidazole.

Scheme I




All the amidino benzimidazoles may be synthesized by these methods in accordance

5 with Scheme II by cyclization of the appropriately substituted aldehyde [(as the bisulphite adduct with diaminobenzamidine] or cyclization of diaminobenzonitrile with the appropriate aldehyde (as the bisulphite adduct) or imidate to give an intermediate nitrile which is then converted to the final product by the Pinner's reaction, i.e. by


10 treating the intermediate nitrile with HCl gas in ethanol to form the imidate followed by reaction with ammonia in ethanol to form the amidine.

Scheme II




- The cyclic amidines (imidazolines etc.) may be synthesized by two routes (Scheme III);
- 5 (A) reaction of the appropriately substituted esters with a diamine in the presence of trimethylaluminium (G. Neef, U. Eder and G. Sauer; J. Org. Chem. 46, 2824 - 2826) and (B) reaction of the appropriately substituted nitrile with ethanolic HCl to give the imidate followed by reaction with the appropriate diamine.

Scheme III




The aldehyde I starting materials may be obtained commercially or may be synthesized
 5 by methods known in the art. For example, the phenoxy derivatives where R_2 or R_3 are
 moieties of the formula:



may be synthesized in one of the following routes (Scheme IV): A.) Reduction of the
 corresponding commercially available nitrile II to the aldehyde with


diisobutylaluminum hydride (A. E. G. Miller, J. W. Biss, and L. H. Schwartzman; J. O. C. 24, 627, 1959), or B.) Nucleophilic displacement reaction of the appropriately substituted halo-aromatic aldehyde or halo-aromatic nitrile with the corresponding phenols. In the second instance the nitriles obtained may either be converted to the final
5 products or reduced with diisobutylaluminum hydride to give the aldehydes.

Scheme IV



Similarly, the piperazine derivatives may be synthesized by nucleophilic displacement
 5 reaction of the appropriately substituted halo-aromatic aldehyde or halo-aromatic nitrile
 with the corresponding piperazine as in Scheme V.

Scheme V



The foregoing reactions are performed in a solvent appropriate to the reagents and materials employed and suitable for the transformation being effected. It is understood by those skilled in the art of organic synthesis that the various

5 functionalities present on the molecule must be consistent with the chemical transformations proposed. This will frequently necessitate judgment as to the order of synthetic steps, protection of reactive groups, and selection of reaction conditions. Reaction conditions compatible with the substituents employed will be apparent to one skilled in the art, as will be the selection of protecting groups where needed.

10

From formula I it is evident that some of the compounds of the invention may have one or more asymmetrical carbon atoms in their structure. It is intended that the present invention include within its scope the stereochemically pure isomeric forms of the compounds as well as their racemates. Stereochemically pure isomeric forms may

15 be obtained by the application of known principles. Enantiomers may be separated from each other by the selective crystallization of the diastomeric salts with optically active bases or by chiral chromatography. Pure stereoisomers may also be prepared

synthetically from appropriate stereochemically pure starting materials, or by using stereospecific reactions.

Suitable pharmaceutical salts are those of inorganic or organic acids, such as
5 hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, nitric acid,
phosphoric acid, acetic acid, succinic acid, oxalic acid, malic acid and the like. Suitable
salts are also those of inorganic or organic bases, such as KOH, NaOH, Ca(OH)₂,
Al(OH)₃, piperidine, morpholine, ethylamine, triethylamine and the like.

10 The pharmaceutically acceptable esters of the present invention are such as would
be readily apparent to those skilled in the art particularly those biolabile esters which are
hydrolyzed under physiological conditions, for example alkoxyalkyl, cycloalkoxyalkyl,
alkenylalkoxyalkyl, aryloxyalkyl, alkoxyaryl, alkylthioalkyl, cycloalkylthioalkyl,
alkenylthioalkyl or alkylthioaryl esters. Particular examples are the pivaloyloxymethyl,
15 acetoxymethyl, phthalidyl, indanyl, and methoxymethyl esters.

Also included within the scope of the invention are the hydrated forms of the
compounds which contain various amounts of water, for instance, the hydrate,
hemihydrate and sesquihydrate forms.

20

The ability of bacteria to quickly respond to changes in the environment is of
utmost importance for their survival. Bacteria are capable of rapidly responding and
adapting to such diverse stimuli as changes in nutrients, osmolarity, temperature, light,
or host environment. These responses may be transient, such as those required for
25 changes in motility or for entry into a host cell. Alternatively, the responses may require
major shifts in gene expression and cell morphology, such as those required for
sporulation, or for survival within a macrophage. The mechanism by which bacteria are
able to sense cues from the physical environment (or from within the cytoplasm) and

process these signals into appropriate responses often involves the so-called "two-component" systems.

As stated above, the antibacterial activity of the compounds of the present invention is based on the inhibition of this "two-component switch" system. All bacteria use this mechanism to control various adaptive/virulence factors to facilitate establishment of a bacterial population in the environment (for example, a bacterial infection in a host). The system invariably consists of a sensor which either activates a kinase or is a part of the kinase, and which upon stimulation, autophosphorylates. This phosphorylated species is a highly active phosphodonor which immediately transfers its phosphate to a "regulatory" component, which in turn initiates the biological response such as transcription or further phosphotransfer in a cascade which eventually ends in regulation of bacterial gene expression. Although each of the kinases and response regulators has a unique sequence (in fact, even functionally identical proteins have slightly different sequences in different species) they share a homologous biochemical mechanism and they share significant homology in the active site.


Accordingly, the present invention provides compounds which exhibit antibiotic activity by inhibiting the autophosphorylation of these bacterial histidine kinases. They also inhibit the transfer of phosphate from phosphorylated histidine kinases to the aspartyl residues of the phosphate acceptor proteins involved in regulation of bacterial gene expression.

The compounds of the present invention have antibacterial activity as determined by the following tests. First, the compounds were tested for their activity in inhibiting the autophosphorylation of Kinase A and the transphosphorylation of Spo0F, two proteins involved in one of the above described signal transduction systems controlling gene expression in bacteria. Representative compounds were then tested for

antibacterial activity against selected organisms by the standard MIC method. The results are set forth below.

Tables 1-6 list examples of compounds of the invention, along with their IC₅₀ values in the HPK *in vitro* assay described below, and MIC value ranges for the selected microorganisms identified below. These examples are merely illustrative of the invention, and are not intended to limit the scope of the claims in any way.

Table 1



Example	R1	R2	R3	R4	IC ₅₀
2	H	H	Ph	H	86
3	H	H	CF ₃	H	169
4	H	H	i-Propyl	H	219
5	H	H	N(Ph) ₂	H	17.4
6	H	OCH ₂ Ph	H	OCH ₂ Ph	15.9
7	H	CF ₃	H	CF ₃	30.8
8	H	H	C(CH ₃) ₂ CH ₂ CH ₃	H	37.6
9	H	t-Butyl	OH	t-Butyl	75.9
10	H	H	t-Butyl	H	70.6
11	H	H	OCH ₂ (4-NO ₂ Ph)	H	19.4
12	H	OC ₁₀ H ₂₁	OC ₁₀ H ₂₁	H	8.3
13	H	H	CHCHPh	H	21.5
14	H	H	N(C ₄ H ₉) ₂	H	35.5
15	H	H	OCH ₂ Ph	H	60.8
16	H	OCH ₂ Ph	OCH ₂ Ph	H	50.7
17	H	H	F	H	480
18	H	I	OH	I	254
19	H	Cl	H	Cl	100
20	H	t-Butyl	H	t-Butyl	19
21	H	Ph	H	Ph	31
22	H	H	NHCH ₂ Ph	H	222
23	H	H	OH	t-Butyl	22
24	H	H	Pyrrolinyl	H	93
25	H	H	N(CH ₂ CH ₂ CN) ₂	H	35.6
26	H	H	N(CH ₂ Ph) ₂	H	17
27	OH	H	H	H	465
28	OCH ₃	H	OCH ₃	Br	66
29	OCH ₃	t-Butyl	H	t-Butyl	76






30	OH	t-Butyl	H	t-Butyl	8.5
31	OH	I	H	I	30.8
32	OH	Br	H	Br	51.1
33	OH	NO ₂	H	NO ₂	74.7
34	OH	t-Butyl	H	H	4
35	OH	H	H	t-Butyl	64
36	OH	Cl	H	Cl	155
37	OH	Br	H	OCH ₃	171
38	OH	H	N(CH ₂ CH ₃) ₂	H	51
39	OH	Ph	H	Ph	23
40	OH	Br	H	NO ₂	199
41	H	Pyridine	H	H	271
42	OCH ₂ COOH	t-Butyl	H	t-Butyl	199
43	H	H		H	40.2
44	H	H		H	24.9
45	H	H		H	28

Table 2




Example	R	R ₁	R ₂	R ₃	R ₄	R ₅	IC ₅₀
46	H	H	H	H	H	CH ₃	44
47	H	H	H	Amidine	H	H	131
48	H	H	H	H	H	H	42
49	H	NO ₂	H	Cl	H	H	29.2
50	H	H	H	NO ₂	H	H	31.7
51	H	H	H	NH ₂	H	H	140.6
52	H	H	H	COOEt	H	H	56
53	H	H	OCH ₃	OCH ₃	OCH ₃	H	28.7
54	H	H	H	OCH ₃	H	H	17.7
55	H	H	OCH ₃	H	OCH ₃	H	59.2
56	H	H	H	t-Butyl	H	H	66.5
57	H	H	NO ₂	NO ₂	H	H	31.6
58	H	H	NH ₂	NH ₂	H	H	31.5
59	H	H	H	Cl	H	H	13.9
60	H	H	Cl	Cl	H	H	11.4
61	NO ₂	H	H	H	H	H	58
62	NH ₂	H	H	H	H	H	19

Table 3



Example	R1	R2	R3	IC ₅₀ (μM)
63	H	H	H	53
64	CF ₃	H	H	48
65	H	CH ₃	H	33
66	Cl	H	Cl	32
67	Cl	Cl	H	32
68	H	t-Butyl	H	31.6
69	H	Cl	H	28
70	H	OCH ₃	H	69.2

Table 4



Example	R	IC ₅₀ (μM)
71		17.3
72		64.3









73		22.0
74		120
75		127
76		85
77		18.5
78		114

Table 5







Example	R ₁	R ₂	R ₃	IC ₅₀
79	H	H		241
80		H		94
81		H		26
82			N(Ph) ₂	22

Table 6



Example	R	R ₁	R ₂	R ₃	R ₄	IC ₅₀
83		H	H	OPh	H	112.4
84		H	H	OPh	H	32.3

85		OH	t-butyl	H	t-butyl	4
86		OH	t-butyl	H	t-butyl	32
87		OH	t-butyl	H	t-butyl	29
88		OH	t-butyl	H	t-butyl	12
89	COOH	OH	t-butyl	H	t-butyl	86
90	NH ₂	OH	t-butyl	H	t-butyl	152
91	CH ₂ NH ₂	OH	t-butyl	H	t-butyl	4

The protocol for the above referenced assay is as follows.

Autophosphorylation of Kinase A and Transphosphorylation of Spo0F Assay

To study the effect of the compounds of the present invention on the signal transduction process in bacteria, the inhibiting effect of the compounds on the sporulation operon proteins Kinase A and Spo0F was examined. Specifically, the inhibition of autophosphorylation of Kinase A and the transphosphorylation of Spo0F was determined in the following assays. The Spo0F response regulator is the primary substrate for phosphorylation by the protein kinase, Kin A, involved in the sporulation process in bacteria. See D. Burbulys, K.A. Trach, J.A. Hoch, *Cell*, **64**, 545-552 (1991). Spo0F and KinA were prepared from recombinant *E. coli* overexpressing the proteins (J. Cavanagh *et al*, *Amino Acids*, **6**, 131-140 (1994) and references therein).

The following stock reagents were either prepared and used promptly or stored at the indicated temperature:

8 X Salts : 2 M KCl (5 mL), 1 M MgCl₂ (800 mL), 1 M CaCl₂ (100 mL), 10 mg/mL phenylmethylsulfonyl fluoride (200 mL), 1 M dithioereitol (50 mL), 0.25 M Na₂EDTA (32 mL) and H₂O 3.82 mL (-20 °C)

5X Loading Dye: 0.5M TRIS-HCl, pH 6.8 (7.5 mL), 10% SDS (2 mL) 0.1% bromophenol blue (0.5 mL), 100% glycerol (3 mL) and 12.5 M 2-mercaptoethanol (0.3 mL)

1-1.3 mg/mL KinA: 15 mM TRIS-HCl, pH 8.0, 6 mM KCl; 4 mM 2-mercaptoethanol; 40% glycerol (-20 °C)

1 mg/mL Spo0F: 17.5 mM TRIS-HCl, pH 8.0; 0.7 mM KCl ; 0.7 mM MgCl₂; 0.7 mM CaCl₂; 5mM 2-mercaptoethanol; 30% glycerol (-20 °C)

5% Stacking Gel: 40% 29:1 acrylamide:bis acrylamide (1.25 mL), 0.5 M TRIS-HCl, pH 6.8 (2.5 mL), 10% SDS (0.1mL), D-H₂O (6.15 mL) 10% ammonium persulfate (100 mL) and TEMED (25 mL)

SDS Running Buffer: TRIS-BASE (3.02 g), glycine (14.4 g),

SDS (1 g), D-H₂O (to 1 L)

The reaction mixture was prepared from 8X Salts (87 µL), 1M TRIS, pH 8 (118 µL), 50% glycerol (63 µL), Spo0F (14.1 µL) and KinA (7.0 µL). Microcentrifuge tubes were charged with the reaction mixture (18.5 µL) and a 1.0 mM solution of the test compound in 5% DMSO (18.5 µL), and incubated for 15 min on ice. 100 mM ATP solution (3.0 µl, containing 625 µCi [³²P]ATP) was added, and the mixture left for 10 minutes at room temperature. The reaction was quenched with 5X loading dye (10 µL per tube) and the samples were loaded on a prepared 5% Stacking Gel, or stored on dry ice until ready for use. The prepared wells were filled with SDS Running Buffer, samples were loaded into the wells, and 80 volts were applied to the gel until the dye front reached the bottom of the stacking gel. The voltage was then increased to 250 volts until electrophoresis was complete. Radioactive bands in the gel

corresponding to phosphorylated KinA and Spo0F were imaged and quantitated with a phosphoimager.

If either enzyme was inhibited (as evidenced by the absence of labelled protein
5 in the developed gel), an IC₅₀ was calculated by running the assay with a range of
inhibitor concentrations from 1 to 500 μ M. After electrophoresis of the reaction
mixtures, percent inhibition was determined by measuring the concentration of
radioactive phosphorus with a phosphoimager and calculating the values using a
software program (BioRad Molecular Analyst).

10

MIC Antimicrobial Assay

The *in vitro* antimicrobial activity of the compounds was determined by the
microdilution broth method following the test method from the National Committee for
Laboratory Standards (NCCLS). This method is described in the NCCLS Document
15 M7-A2, Vol.10, No.8 "Methods for Dilution Antimicrobial Susceptibility Test for
Bacteria that Grow Aerobically - Second Edition."

In this method two-fold serial dilutions of drug in cation supplemented Mueller-
Hinton broth are added to wells in microdilution trays. The test organisms are prepared
20 by adjusting the turbidity of actively growing broth cultures so that the final
concentration of test organism after it is added to the wells is approximately 5×10^4
CFUs/well).

Following inoculation of the microdilution trays, the trays are incubated at 35°C
25 for 16-20 hours and then read. The MIC is the lowest concentration of test compound
that completely inhibits growth of the test organism. The amount of growth in the wells
containing the test compound is compared with the amount of growth in the growth-
control wells (no test compound) used in each tray.

The antibacterial results demonstrated that the compounds according to the invention have a broad spectrum of activity against the bacterial strains tested including one or more of the following test organisms:

- 5 *S. aureus* (ATCC 29213)
 Methicillin resistant *S. aureus* (oc 2089, 2878 and 667)
 E. faecalis (oc 3041)
 E. faecium (oc 2993)
 E. coli (oc 2530)
- 10 *S. epidermidis* (oc 2603)
 K. pneumoniae (oc 1943)
 P. aeruginosa (oc 161)

The compounds according to this invention which have good antimicrobial activity can
15 be used in antimicrobial compositions, as antiseptics or disinfectants for local and general application, topically for prevention or treatment of skin lesions, and, where in vivo activity is present, for systemic treatment of bacterial infections in the form of conventional antimicrobial pharmaceutical compositions appropriate for the intended use.

20

As antiseptics for human or veterinary use, the concentration of active compound can vary from 0.01% to 5% by weight according to the use and the chosen formulation. Thus, it is possible to prepare foaming detergent solutions to be used by surgeons and nursing staff for washing their hands or to be used for cleansing
25 dermatological lesions such as impetigo, pityriasis and leg ulcers. Foaming detergent solutions are also used as shampoos or for the preparation of shower gels and foaming lotions. Foaming solutions containing the compounds according to the invention are obtained using amphoteric, anionic, cationic, or non-ionic surfactants at a concentration of 0.3 to 30%, humectants such as glycols and polyethylene glycols at a concentration

of 0 to 20%, ethylene oxide and polypropylene copolymers at a concentration of 0 to 20%, and an alcohol (ethanol, isopropyl or benzyl alcohol) or a polyol, such as glycerol, at a concentration of 0 to 15%, as well as agents for complexing Ca^{++} , Mg^{++} and heavy metal ions, salts for providing an appropriate buffer capacity, agents for
5 imparting viscosity, natural, cellulosic or synthetic polymers, fragrances, preservatives and colorants.

If the compound according to the invention has a poor solubility in water, it is possible to use microemulsions, micellar solutions or similar surfactant systems which
10 permit solubilization in water. These solutions can be used in diluted or undiluted form and can be dispensed by means of a vasopump or liquified or non-liquified propellants.

With the same constituents at appropriate concentrations, the compounds according to the invention can also be used to prepare simple aqueous solutions or
15 sprays for making operative fields antiseptic, for postoperative treatments, for the treatment of burns, superinfected eczema, gluteal erythema, wounds or acne, or for deodorants. Alternatively, the compounds can be formulated in simple alcoholic solutions in the form of sprays containing 20 to 80% by weight of alcohol, optionally including excipients which make it possible to penetrate the keratinized layers of the
20 skin and superficial body growths, such as Azone. These solutions can be used for making the skin antiseptic before puncture, for preparing the operative field, by nursing staff for making their hands antiseptic and for treating closed infected dermatosis, folliculitis, perionchya and acne. Compositions for topical application may take the form of liquids, creams or gels, containing a therapeutically effective concentration of a
25 compound of the invention admixed with a dermatologically acceptable carrier.

Moreover, the rapid lethal action on germs of the compounds of the invention enables them to be used as surface disinfectants at concentrations which can vary from 0.1 to 4% by weight. In this case, the product is used in preparations such as aqueous

or nonaqueous foaming detergent solutions, sprays or nebulizers. Preparations of this type are particularly useful in the hospital or veterinary sectors or agrifoodstuff industries. These preparations can contain the same constituents as those used in the antiseptic formulations, although a wide variety of organic solvents may be added.

5

Finally, for those compounds which exhibit in vivo activity, the invention further provides a method of treating bacterial infections, or enhancing the activity of other antibacterial agents, in warm-blooded animals, which comprises administering to the animals a compound of the invention alone or in admixture with a diluent or in the form of a medicament according to the invention.

10

When the compounds are employed for the above utility, they may be combined with one or more pharmaceutically acceptable carriers, e.g., solvents, diluents, and the like, and may be administered orally in such forms as tablets, capsules, dispersible powders, granules, or suspensions containing for example, from about 0.5% to 5% of suspending agent, syrups containing, for example, from about 10% to 50% of sugar, and elixirs containing, for example, from about 20% to 50% ethanol, and the like, or parenterally in the form of sterile injectable solutions or suspensions containing from about 0.5% to 5% suspending agent in an isotonic medium. These pharmaceutical preparations may contain, for example, from about 0.5% up to about 90% of the active ingredient in combination with the carrier, more usually between 5% and 60% by weight.

15

20

In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed. Solid carriers include starch, lactose, dicalcium phosphate, microcrystalline cellulose, sucrose and kaolin, while liquid carriers include sterile water, polyethylene glycols, non-ionic surfactants and edible oils such as corn, peanut and sesame oils, as are appropriate to the nature of the active ingredient and the particular form of administration desired. Adjuvants customarily

25

employed in the preparation of pharmaceutical compositions may be advantageously included, such as flavoring agents, coloring agents, preserving agents, and antioxidants, for example, vitamin E, ascorbic acid, BHT and BHA.

- 5 The preferred pharmaceutical compositions from the standpoint of ease of preparation and administration are solid compositions, particularly tablets and hard-filled or liquid-filled capsules. Oral administration of the compounds is preferred.

10 These active compounds may also be administered parenterally or intraperitoneally. Solutions or suspensions of these active compounds as a free base or pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxypropyl-cellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof in oils. Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent
15 the growth of microorganisms.

20 The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol and liquid polyethylene glycol), suitable mixtures thereof, and
25 vegetable oils.

The effective dosage of active ingredient employed may vary depending on the particular compound employed, the mode of administration and the severity of the condition being treated. However, in general, satisfactory results are obtained when the

compounds of the invention are administered at a daily dosage of from about 0.1 mg/kg to about 400 mg/kg of animal body weight, preferably given in divided doses two to four times a day, or in sustained release form. For most large mammals the total daily dosage is from about 0.07 g to 7.0 g, preferably from about 100 mg to 1000 mg

5 Dosage forms suitable for internal use comprise from about 100 mg to 500 mg of the active compound in intimate admixture with a solid or liquid pharmaceutically acceptable carrier. This dosage regimen may be adjusted to provide the optimal therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic

10 situation.

The production of the above-mentioned pharmaceutical compositions and medicaments is carried out by any method known in the art, for example, by mixing the active ingredient(s) with the diluent(s) to form a pharmaceutical composition (e.g. a

15 granulate) and then forming the composition into the medicament (e.g. tablets).

The following examples describe in detail the chemical synthesis of representative compounds of the present invention. The procedures are illustrations, and the invention should not be construed as being limited by chemical reactions and conditions they

20 express. No attempt has been made to optimize the yields obtained in these reactions, and it would be obvious to one skilled in the art that variations in reaction times, temperatures, solvents, and/or reagents could increase the yields.

EXAMPLE 1

(ROUTE A)

2-[4-(1,1-dimethylethyl)phenyl]-1H-Benzimidazole-5-carboximidamide

(a) 3,4-Diaminobenzimidate

30 To 500 mL of ethanol in an ice bath was bubbled HCl (g) until the ethanol was saturated with the gas. 3,4-Diaminobenzonitrile (40.0 g, 0.300 mol) was dissolved in 500 mL of CH₂Cl₂ and added to the cooled ethanol solution. HCl (g) was continued to

be bubbled through the solution at 0 °C. The reaction was sealed and warmed up to 25 °C. The reaction was followed by TLC along with IR spectroscopy for the disappearance of the nitrile. Once there were no more evidence of the nitrile, the reaction mixture was put under *vacuo* to remove 75% of the solvents. Ether (500 mL) was added, the precipitate was filtered, washed with ether (3 x 250 mL), and dried under nitrogen. The 3,4-diaminobenzimidate was used to synthesize 3,4-diaminobenzamidine dihydrochloride without any further purification.

10 (b) 3,4-Diaminobenzamidine dihydrochloride


To 500 mL of ethanol at 25 °C was bubbled NH₃ (g) until the ethanol was saturated with the gas. The imidate was added and the mixture was heated to 80 °C under nitrogen for 4 h. The reaction was followed by TLC with 1:1 Hex/EtOAc. Once the reaction was over, the reaction solution was place under *vacuo* to remove all solvents. 15 The residue was recrystallized from 2N HCl to afford 3,4-diaminobenzamidine dihydrochloride (51.4 g, 76%) as an off-white solid: ¹H NMR (DMSO-*d*₆) δ 9.14 (bs, 2 NH), 8.92 (bs, 2 NH), 7.59 (s, 1 arom-H), 7.51 (d, 1 arom-H, J = 8Hz), 6.95 (d, 1 arom-H, J = 8Hz).

20 (c) 2-[4-(1,1-dimethylethyl)phenyl]-1H-Benzimidazole-5-carboximidamide

To 4-tert-butylbenzaldehyde (0.4000 g, 0.00247 mol) was added 3 mL of 40% NaHSO₃ (aq). The mixture was stirred for 2 h at 25 °C. 3,4-Diaminobenzamidine dihydrochloride (0.56 g, 0.0025 mol) and ethanol (8 mL) were added. The mixture was 25 refluxed for 3 h. The reaction was then poured into water. The precipitate filtered and recrystallized from 2N HCl to afford 1H-benzimidazole-5-carboximidamide, 2-[4-(1,1-dimethylethyl)phenyl]- (400 mg, 44%) as a white solid: ¹H NMR (DMSO-*d*₆) δ 9.47 (bs, 2 NH), 9.15 (bs, 2 NH), 8.30 (d, 2 arom-H, J = 8Hz), 8.21 (s, 1 arom-H), 7.88 (d, 1 arom-H, J = 8Hz), 7.73 (d, 1 arom-H, J = 9Hz), 7.68 (d, 2 30 arom-H, J = 8Hz), 1.35 (s, 9 tert-butyl-H).

The following compounds are prepared using the procedure of Example 1 (Route A) and the appropriate starting materials. The starting materials are commercially available or may be prepared in accordance with the following Reference Examples.

Table 7



Example	R1	R2	R3	R4	m.p.
2	H	H	Ph	H	297-298
3	H	H	CF ₃	H	268-270
4	H	H	i-Propyl	H	282-283
5	H	H	N(Ph) ₂	H	>300
6	H	OCH ₂ Ph	H	OCH ₂ Ph	195-197
7	H	CF ₃	H	CF ₃	199-201
8	H	H	C(CH ₃) ₂ CH ₂ CH ₃	H	237-238
9	H	t-Butyl	OH	t-Butyl	286-287
10	H	H	t-Butyl	H	249-251
11	H	H	OCH ₂ (4-NO ₂ Ph)	H	290-291
12	H	OC ₁₀ H ₂₁	OC ₁₀ H ₂₁	H	255-257
13	H	H	CHCHPh	H	>300
14	H	H	N(C ₄ H ₉) ₂	H	291-292
15	H	H	OCH ₂ Ph	H	253-254
16	H	OCH ₂ Ph	OCH ₂ Ph	H	247-248
17	H	H	F	H	>300
18	H	I	OH	I	>300
19	H	Cl	H	Cl	>300
20	H	t-Butyl	H	t-Butyl	270-275
21	H	Ph	H	Ph	204-206
22	H	H	NHCH ₂ Ph	H	265-267
23	H	H	OH	t-Butyl	299-300
24	H	H	Pyrolinyl	H	>300
25	H	H	N(CH ₂ CH ₂ CN) ₂	H	284-286
26	H	H	N(CH ₂ Ph) ₂	H	271-273
27	OH	H	H	H	>300
28	OCH ₃	H	OCH ₃	Br	254-255
29	OCH ₃	t-Butyl	H	t-Butyl	>250
30	OH	t-Butyl	H	t-Butyl	272-273
31	OH	I	H	I	223-224
32	OH	Br	H	Br	262-263
33	OH	NO ₂	H	NO ₂	>300
34	OH	t-Butyl	H	H	240-246
35	OH	H	H	t-Butyl	>265





36	OH	Cl	H	Cl	>300
37	OH	Br	H	OCH ₃	249-250
38	OH	H	N(CH ₂ CH ₃) ₂	H	>250
39	OH	Ph	H	Ph	>300
40	OH	Br	H	NO ₂	>300
41	H	Pyridine	H	H	230-236
42	OCH ₂ COOH	t-Butyl	H	t-Butyl	>300
43	H	H		H	>300
44	H	H		H	>300
45	H	H		H	237-238


Table 8



Example	R	R ₁	R ₂	R ₃	R ₄	R ₅	m.p.
46	H	H	H	H	H	CH ₃	>245
47	H	H	H	Amidine	H	H	255-257
48	H	H	H	H	H	H	253
49	H	NO ₂	H	Cl	H	H	296-298
50	H	H	H	NO ₂	H	H	298-300
51	H	H	H	NH ₂	H	H	>300
52	H	H	H	COOEt	H	H	261-262
53	H	H	OCH ₃	OCH ₃	OCH ₃	H	286-288
54	H	H	H	OCH ₃	H	H	269-270
55	H	H	OCH ₃	H	OCH ₃	H	236-238
56	H	H	H	t-Butyl	H	H	291-292
57	H	H	NO ₂	NO ₂	H	H	234-236
58	H	H	NH ₂	NH ₂	H	H	282-283


59	H	H	H	Cl	H	H	>300
60	H	H	Cl	Cl	H	H	>300
61	NO ₂	H	H	H	H	H	173-176
62	NH ₂	H	H	H	H	H	168-170

Table 9




Example	R ₁	R ₂	R ₃	m.p.
63	H	H	H	173-174
64	CF ₃	H	H	257-259
65	H	CH ₃	H	>300
66	Cl	H	Cl	292-293
67	Cl	Cl	H	273-274
68	H	t-Butyl	H	265-266
69	H	Cl	H	294-296
70	H	OCH ₃	H	161-163

Table 10




Example	R	m.p.
71		>300
72		>300

73




297-298

74




>300

75




297-298

76




291

77




78




196

Table 11







Example	R ₁	R ₂	R ₃	m.p.
79	H	H		213-215
80		H		245-248
81		H		>300
82			N(Ph) ₂	283-284

Table 12



Example	R	R ₁	R ₂	R ₃	R ₄	m.p.
83		H	H	OPh	H	257-258
84		H	H	OPh	H	>300

85		OH	t-butyl	H	t-butyl	275-276
86		OH	t-butyl	H	t-butyl	>300
87		OH	t-butyl	H	t-butyl	>300
88		OH	t-butyl	H	t-butyl	>260
89	COOH	OH	t-butyl	H	t-butyl	217-219

EXAMPLE 93

ROUTE B(i)

2-(4-Phenoxyphenyl)-1H-benzimidazole-5-carboximidamide

5 (a) 2-(4-phenoxyphenyl)-1H-Benzimidazole-5-nitrile

To 4-phenoxybenzaldehyde (4.0 g, 0.020 mol) was added 20 mL of 40% NaHSO₃ (aq). The mixture was stirred for 2 h at 25 °C. 3,4-Diaminobenzonitrile (3.36 g, 0.0252 mol) and ethanol (50 mL) were added. The mixture was refluxed for 4 h. The reaction was then poured into water. The precipitate was filtered and recrystallized from ethyl acetate to afford 3.6 g (57%) of 2-(4-phenoxyphenyl)-1H-benzimidazole-5-nitrile.

(b) Imidate

To 50 mL of ethanol in an ice bath was bubbled HCl(g) until the ethanol was saturated with the gas. 2-(4-phenoxyphenyl)-1H-benzimidazole-5-nitrile (0.50 g, 0.0016 mol)

was dissolved in 50 mL of CH₂Cl₂ and added to the cooled ethanol solution. HCl (g) was continued to be bubbled through the solution at 0 °C. The reaction was sealed and warmed up to 25 °C. The reaction was followed by TLC along with IR spectroscopy for the disappearance of the nitrile. Once there were no more evidence of the nitrile, the reaction mixture was put under *vacuo* to remove 75% of the solvents. Ether (50 mL) was added, the precipitate was filtered, washed with ether (3 x 50 mL), and dried under nitrogen. The product was used without any further purification to make the amidine dihydrochloride.

**(c) 2-(4-phenoxyphenyl)-1H-benzimidazole-5-carboximidamide,
10 dihydrochloride**

To 100 mL of ethanol at 25°C was bubbled NH₃ (g) until the ethanol was saturated with the gas. The imidate was added and the mixture was heated to 70°C under nitrogen for 4 h. The reaction was followed by TLC with 1:1 Hex/EtOAc. Once the reaction was over, the reaction solution was placed under *vacuo* to remove all solvents.

15 The residue was recrystallized from 2N HCl to afford the amidine 2-(4-phenoxyphenyl)-1H-benzimidazole-5-carboximidamide (350 mg, 54%) as a white solid: ¹H NMR (DMSO-_d₆) δ 9.50 (bs, 2 NH), 9.18 (bs, 2 NH), 8.41 (d, 2 arom-H, J = 9Hz), 8.22 (s, 1 arom-H), 7.89 (d, 1 arom-H, J = 8Hz), 7.80 (d, 1 arom-H, J = 8Hz), 7.51 (d, 1 arom-H, J = 8Hz), 7.48 (d, 1 arom-H, J = 8Hz), 7.18 (m, 5
20 arom-H).

EXAMPLE 94

ROUTE B(ii)

2-(4-hydroxyphenyl)-1H-Benzimidazole-5-nitrile

25

To 50 mL of ethanol in an ice bath was bubbled HCl (g) until the ethanol was saturated with the gas. 4-Hydroxybenzonitrile (2.7 g, 0.023 mol) was dissolved in 50 mL of CH₂Cl₂ and added to the cooled ethanol solution. HCl (g) was continued to be bubbled

through the solution at 0 °C. The reaction was sealed and warmed up to 25 °C. The reaction was followed by TLC along with IR spectroscopy for the disappearance of the nitrile. Once there were no more evidence of the nitrile, the reaction mixture was put under *vacuo* to remove 75% of the solvents. Ether (50mL) was added, the precipitate
5 filtered, washed with ether (3 x 50mL) and dried under nitrogen. The imide was used to make 2-(4-hydroxyphenyl)-1H-benzimidazole-5-nitrile without any further purification.

A mixture of 3,4-diaminobenzonitrile (3.0 g, 0.023 mol) and the imide in 500 mL of
10 methanol was heated at 70°C under nitrogen. The reaction was followed by TLC with 1:1 Hex/EtOAc. After 4 h the reaction solution was placed under *vacuo* to remove all solvents. The residue was recrystallized from ethyl acetate to afford 4.5 g (85%) of 2-(4-hydroxyphenyl)-1H-benzimidazole-5-nitrile.

15

EXAMPLE 95

5-(4,5-Dihydro-1H-imidazol-2-yl)-2-(4-phenoxyphenyl)-1H-
Benzimidazole

(a) 2-(4-phenoxyphenyl)-1H-Benzimidazole-5-carboxylic acid, methyl
20 ester

To 4-phenoxybenzaldehyde (3.0 g, 0.02 mol) was added 21 mL of 40% NaHSO₃ (aq). The mixture was stirred for 1 h at 25 °C. Methyl-3,4-diaminobenzoate (2.5 g, 0.02 mol) and ethanol (15 mL) were added. The mixture was refluxed for 4 h. The reaction was then poured into water. The precipitate filtered and recrystallized from EtOAc to
25 afford 2-(4-phenoxyphenyl)-1H-benzimidazole-5-carboxylic acid, methyl ester (5.06 g, 97%) as a white solid.

(b) 5-(4,5-dihydro-1H-imidazol-2-yl)-2-(4-phenoxyphenyl)-1H-benzimidazole

Ethylenediamine (0.70 g, 0.0116 mol) in benzene (10mL) was added dropwise to a stirred solution of trimethyl aluminum (2.0M, 0.12 mol) such that the temperature does not exceed 10 °C. At the end of methane evolution, the ester 2-(4-phenoxyphenyl)-1H-Benzimidazole-5-carboxylic acid, methyl ester (2.5 g, 0.00726 mol) was gradually added at 25 °C. The reaction mixture was then refluxed for 3 h. The reaction was cooled and treated with 3 mL of H₂O, diluted with MeOH, 10 mL of CH₂Cl₂ and refluxed for 15 min. The mixture was filtered and washed with EtOAc to remove aluminum hydroxides. The filtrate was concentrated and the residue was recrystallized from 2N HCl to afford 5-(4,5-dihydro-1H-imidazol-2-yl)-2-(4-phenoxyphenyl)-1H-benzimidazole (2.3g, 74%) as a white solid.

EXAMPLE 96

2-[5-(4,5-dihydro-4,4-dimethyl-1H-imidazol-2-yl)-1H-benzimidazol-2-yl]-3,5-bis(1,1-dimethylethyl)-phenol

15

(a) 2-[3,5-bis(1,1-dimethylethyl)-2-hydroxyphenyl]-1H-Benzimidazole-5-carbonitrile

To 3,5-di-tert-butylsalicylaldehyde (8.98 g, 0.0383 mol) was added 55 mL of 40% NaHSO₃ (aq). The mixture was stirred for 5 h at 25 °C. 3,4-Diaminobenzonitrile (5.13 g, 0.0385 mol) and ethanol (300 mL) was added. The mixture was refluxed for 24 h. The reaction was then poured into water. The precipitate was filtered and recrystallized from CH₂Cl₂/Hexanes to afford 1H-benzimidazole-5-carbonitrile, 2-[3,5-bis(1,1-dimethylethyl)-2-hydroxyphenyl]- (10.86g, 82%) as a white solid.

(b) Phenol, 2-[5-(4,5-dihydro-4,4-dimethyl-1H-imidazol-2-yl)-1H-benzimidazol-2-yl]-3,5-bis(1,1-dimethylethyl)-

To 60 mL of ethanol in an ice bath was bubbled HCl (g) until the ethanol was saturated with the gas. 1H-benzimidazole-5-carbonitrile, 2-[3,5-bis(1,1-dimethylethyl)-2-

- hydroxyphenyl]- (0.5236 g, 1.508 mmol) was added to the cooled ethanol solution. HCl (g) was continued to be bubbled through the solution at 0 °C. The reaction was sealed and warmed up to 25 °C. The reaction was followed by TLC along with IR spectroscopy for the disappearance of the nitrile. The reaction mixture was then put
- 5 under *vacuo* to remove 75% of the solvents. Ether (100 mL) was added and the precipitate was filtered, washed with ether (3 x 250 mL) and dried under nitrogen to yield the imidate. The imidate was used to make the amidine without any further purification.
- 10 To the imidate in MeOH (10 mL) was added 1,2-diamino-2-methylpropane (0.19 mL, 0.0018 mol). The mixture refluxed for 16 h under N₂. After cooling, the reaction solution was placed under *vacuo* to remove all solvents. The residue was recrystallized from 2N HCl to afford 0.400 g (63%) of phenol, 2-[5-(4,5-dihydro-4,4-
- 15 dimethyl-1H-imidazol-2-yl)-1H-benzimidazol-2-yl]-3,5-bis(1,1-dimethylethyl)- as a white solid.

EXAMPLE 97

7-amino-2-[4-(diphenylamino)phenyl]-1H-benzimidazole-5-
carboximidamide

20

(a) 4-Amino-3,5-dinitrobenzonitrile

To 4-chloro-3,5-dinitrobenzonitrile (9.8 g, 0.043 mol) in EtOAc (30 mL) and DMSO (20 mL) was added 1M ethanolic ammonia (100 mL). The mixture was stirred for 1 h and filtered to afford the 4-amino-3,5-dinitrobenzonitrile (8.78 g, 98%) as a yellow solid.

25 (b) 3,4-Diamino-5-nitrobenzonitrile

To 4-amino-3,5-dinitrobenzonitrile (9.4 g, 0.045 mol) in EtOH (100 mL) was added SnCl₂ (30 g, 0.16 mol). The mixture was refluxed for 1 h. The reaction was then filtered and the filtrate concentrated to afford 3,4-diamino-5-nitrobenzonitrile (3.4 g, 42%) of an orange solid.

(c) 3,4-Diamino-5-nitrobenzimidate:

To 100 mL of ethanol in an ice bath was bubbled HCl (g) until the ethanol was saturated with the gas. 3,4-Diamino-5-nitrobenzonitrile (2.0 g, 0.011 mol) was dissolved in 20 mL of CH₂Cl₂ and added to the cooled ethanol solution. HCl (g) was continued to be bubbled through the solution at 0 °C. The reaction was sealed and warmed up to 25 °C. The reaction was followed by TLC along with IR spectroscopy for the disappearance of the nitrile. Once there were no more evidence of the nitrile, the reaction mixture was put under *vacuo* to remove 75% of the solvents. Ether (100 mL) was added, the precipitate was filtered, washed with ether (3 x 50 mL), and dried under nitrogen. The 3,4-diaminobenzimidate was used to synthesize 3,4-diamino-5-nitrobenzamidine dihydrochloride without any further purification.

(d) 3,4-Diamino-5-nitrobenzamidine dihydrochloride:

To 500 mL of ethanol at 25 °C was bubbled NH₃ (g) until the ethanol was saturated with the gas. The imidate was added and the mixture was heated to 80 °C under nitrogen for 4 h. The reaction was followed by TLC with 1:1 Hex/EtOAc. Once the reaction was over, the reaction solution was placed under *vacuo* to remove all solvents. The residue was recrystallized from 2N HCl to afford 3,4-diamino-5-nitrobenzamidine dihydrochloride (1.86 g, 62%) as a rusty color solid.

(e) 7-nitro-2-[4-(diphenylamino)phenyl]-1H-Benzimidazole-5-carboximidamide

To 4-phenoxybenzaldehyde (430 mg, 2.17 mmol) was added 2 mL of 40% NaHSO₃ (aq). The mixture was stirred for 0.75 h at 25 °C. 3,4-Diamino-5-nitrobenzonitrile dihydrochloride (402 mg, 1.50 mmol) and ethanol (5 mL) were added. The mixture was refluxed for 2.5 h. The reaction was then poured into water. The precipitate was filtered and recrystallized from ethyl acetate to afford (79 mg, 12%) of 1H-benzimidazole-5-carboximidamide, 7-nitro-2-[4-(diphenylamino)phenyl]- as a light yellow solid.

(f) **7-amino-2-[4-(diphenylamino)phenyl]-1H-Benzimidazole-5-carboximidamide**

A mixture of 1H-benzimidazole-5-carboximidamide, 7-nitro-2-[4-(diphenylamino)phenyl]- (50 mg, 0.11 mmol), catalytic amounts of 10% Pd/C and HCl (12N) in EtOH (20 mL) was placed in a Parr reaction bottle under H₂ (50 psi) at 25 °C with shaking for 2 h. The reaction was then filtered over celite and the filtrate concentrated. The residue was recrystallized from 2N HCl/EtOH to afford **7-amino-2-[4-(diphenylamino)phenyl]-1H-benzimidazole-5-carboximidamide**, (42 mg, 92%) as a white solid.

10

Reference Example 1

4-(4-Methoxyphenoxy)benzaldehyde

A mixture of 4-methoxyphenol (1.0 g, 0.00806 mol), 4-bromobenzaldehyde (1.49 g, 0.00805 mol), CuO (0.64 g, 0.00805 mol), potassium carbonate (1.11 g, 0.00803 mol) in pyridine (100 mL) was refluxed for 60 h. The reaction mixture was filtered into ice/water and the product was extracted with ethyl acetate. The organic layer was washed with water, dried (MgSO₄), concentrated and chromatographed with CH₂Cl₂ to afford 1.56 g (85%) of **4-(4-methoxyphenoxy)benzaldehyde**.

20

Reference Example 2

2-[4-(4-cyanophenoxy)phenyl]-1H-Benzimidazole-5-carbonitrile

A mixture of 1H-Benzimidazole-5-nitrile, 2-(4-hydroxyphenyl)- (2.6 g, 0.011 mol), potassium carbonate (4.0 g, 0.029 mol), and 4-fluorobenzonitrile (1.2 g, 0.0099 mol) in DMF (100 mL) was refluxed for 18 h. The mixture was then filtered (to remove the potassium carbonate) into an ice/water bath. The precipitate was collected and recrystallized from ethyl acetate to afford 0.7 g (21%) of **2-[4-(4-cyanophenoxy)phenyl]-1H-benzimidazole-5-carbonitrile**.

Reference Example 3

4-(4-*tert*-Butylphenoxy)benzaldehyde

- 5 A mixture of 4-fluorobenzaldehyde (2.0 g, 0.0161 mol), 4-*tert*-butylphenol (2.42 g, 0.161 mol), and K₂CO₃ (9.0 g, 0.065 mol) in DMF (50 mL) was heated at 80 °C. After 6 h, the reaction mixture was filtered into ice/water and the product was extracted with ethyl acetate. The organic layer was washed with water, dried (MgSO₄), concentrated and chromatographed with 1:1 hexane/ethyl acetate to give 3.9 g (95%) of
- 10 **4-(4-*tert*-butylphenoxy)benzaldehyde.**

Reference Example 4

4-(4-Phenyl-1-piperazinyl)-benzaldehyde

- 15 A solution of 4-fluorobenzaldehyde (0.86 mL, 8.06 mmol), 1-phenylpiperazine (4.89 mL, 32 mmol), and potassium carbonate (1.10g, 8.06 mmol) in acetonitrile was warmed to reflux under N₂. After 60h, the reaction mixture was allowed to cool to 25 °C, diluted with ethyl acetate (50 mL), washed with water (2 x 100 mL), filtered through a plug of silica, and was concentrated *in vacuo* to afford a yellow residue. The
- 20 residue was washed with warm methanol, and dried under vacuum to afford **benzaldehyde, 4-(4-phenyl-1-piperazinyl)-** (1.71g, 79%) was a white solid: ¹H NMR(CDCl₃) δ 9.82 (s, 1H), 7.90 (d, J = 8.8 Hz, 2H), 7.33-7.20 (m, 3H), 7.01-6.80 (m, 4H), 3.38-3.28 (m, 4H), 3.20-3.13 (m, 4H).
- 25 solid: ¹H NMR(CDCl₃) δ 9.78 (s, 1H), 8.54 (s, 1H), 7.89 (d, J = 7.4 Hz, 1H), 6.83 (d, J = 7.4 Hz, 1H), 3.75 (m, 4H), 1.80-1.65 (m, 6H).

Reference Example 5

3-*tert*-Butyl-2-hydroxybenzaldehyde

A mixture of 2-*tert*-butylphenol (15 g, 0.010 mol), powdered NaOH (20 g, 0.50 mol),
5 H₂O (3.6 mL), and CHCl₃ (100 mL) was heated at 50 °C for 1 h. Then more
powdered NaOH (12 g, 0.30 mol) was added at an increment of 2 g every 15 min. At
the end of the NaOH additions, the reaction was continued to be heated at 56 °C for 1
h. After cooling, the reaction was acidified with HCl (12N). The chloroform fraction
was separated, dried (MgSO₄), and concentrated. The residue was loaded on a silica
10 gel column eluted with CH₂Cl₂ to give 3-*tert*-butyl-2-hydroxybenzaldehyde (4.6
g, 26%) as a solid.

Reference Example 6

2-(*tert*-Butylhydroxyacetate)-3,5-di-*tert*-butylbenzaldehyde

15 A solution of 3,5-di-*tert*-butyl-2-hydroxybenzaldehyde (1g, 4.27 mmol) and sodium
hydride (144 mg, 6.0 mmol) in dimethylformamide (50 mL) at 25 °C under N₂ was
treated with *tert*-butyl bromoacetate (0.75 mL, 4.7 mmol) via syringe. After 8h, the
reaction mixture diluted with ethyl acetate (50 mL), washed with water (2 x 25 mL),
20 filtered through a plug of silica and was concentrated *in vacuo* to afford 2-(*tert*-butyl
hydroxyacetate)-3,5-di-*tert*-butylbenzaldehyde (1.48 g, 100%) as a yellow
residue that was used without further purification: ¹H NMR(CDCl₃) δ 10.35 (s, 1H),
7.65 (d, J = 3.2 Hz, 1H), 7.62 (d, J = 3.2 Hz, 1H), 4.42 (s, 1H), 1.56 (s, 9H), 1.45
(s, 9H), 1.37 (s, 9H).

25


Reference Example 13

3,5-Diphenylbenzaldehyde

A solution of 3,5-diphenylbenzoic acid (1 g, 3.65 mmol) in tetrahydrofuran (50 mL) was treated with a solution of lithium aluminum hydride (2.6 mL, 1M, 2.6 mmol) in tetrahydrofuran at 25 °C under N₂. After 3 h, the reaction mixture was carefully treated with satd. aqueous ammonium chloride (1 mL), diluted with ethyl acetate (50 mL),
5 washed with satd. aqueous ammonium chloride (2 x 50 mL), dried (MgSO₄), and was concentrated *in vacuo* to afford a white solid that was used without any further purification. This solid residue was diluted with ethyl ether (20 mL) was treated with manganese dioxide (6.6 g, 76.7 mmol) at 25 °C. After 12h, the reaction mixture was filtered through a plug of celite, and was concentrated *in vacuo*. Purification by flash
10 chromatography afforded **2** (410 mg, 43%) as a solid: ¹H NMR(CDCl₃) δ 10.23 (s, 1H), 8.25-8.20 (m, 5H), 8.14 (s, 2H), 7.66-7.45 (m, 6H).

CLAIMS

1. A compound having antimicrobial activity selected from those of the Formula I :




Formula I

wherein:

R₁ is selected from H, OH, and (C₁-C₆)alkoxy;

R₂, R₃ and R₄ are independently selected from H, OH, (C₁-C₁₀)alkyl (branched or
 10 unbranched), (C₁-C₁₀)alkoxy, -O(C₁-C₆)alkylphenyl, -OCH₂(4-NO₂phenyl), CF₃,
 halo, phenyl, substituted phenyl (wherein the substituents are selected from halogen,
 (C₁-C₆)alkoxy, nitro, NH₂, and (C₁-C₆)alkyl), -(C₁-C₆)alkylphenyl, nitro, -
 O(CH₂)_qN((C₁-C₆)alkyl)₂ wherein q is 1-3, -NR₁₇R₁₈, 2-thiazole optionally
 substituted by a (C₁-C₆) alkyl group, -CH=CH-phenyl, -O-piperazine, pyrrolinyl, and


15 a moiety of the formula:



wherein n is an integer from 0-2;

R₁₂, R₁₃, R₁₄, R₁₅ and R₁₆ are independently selected from H, nitro, halo,
 CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and amino, amidine, sulfonate,
 20 -C(O)O(C₁-C₆)alkyl, and carboxyl;

or R₁₃ and R₁₄ or R₁₄ and R₁₅ taken together form a fused 5 membered
 heterocyclic ring of the formula:




which may be optionally substituted wherein the substituent is selected from amino and methyl;

R₁₇ and R₁₈ are independently selected from H, benzyl, phenyl, (C₁-C₆)alkyl,

5 CH₂CN, CH₂CH₂CN, or R₁₇ and R₁₈ taken together with the nitrogen to which they are attached form a 5-6 membered monocyclic heterocyclic ring optionally containing 1-2 additional nitrogen atoms;


or R₂ and R₃ or R₃ and R₄ taken together with the phenyl ring to which they are attached form a moiety selected from those of the formulae:



is CH, O or N and R₂₀ is (C₁-C₃)alkyl or OH;

R₅ is selected from H, amino and amidino;

R₆ is selected from nitro and moieties of the formula:



15


and



wherein:

R₉ and R₁₀ are H or R₉ and R₁₀ taken together form a 5, 6 or 7 membered monocyclic heterocyclic ring optionally containing 1 or 2 additional nitrogens and optionally substituted wherein the substituent is dimethyl, or R₉ and R₁₀ taken together form a bicyclic heterocyclic ring of the formula;

20



R₇ is selected from H, amino and nitro;

R₈ is selected from H and methyl;

and the pharmaceutically acceptable salts, esters and prodrug forms thereof.


5

2. A compound having antimicrobial activity according to claim 1 wherein:

R₁ is selected from H, OH, (C₁-C₆)alkoxy;

R₂ is selected from H, (C₁-C₁₀)alkyl (branched or unbranched), (C₁-C₁₀)alkoxy; -

O(C₁-C₁₀)alkylphenyl, CF₃, halo, phenyl, nitro, pyridyl, and a moiety of the formula:




10

R₃ is selected from H, OH, (C₁-C₁₀)alkyl (branched or unbranched), (C₁-C₁₀)alkoxy,

-O(C₁-C₆)alkylphenyl, -OCH₂(4-NO₂phenyl), CF₃, halo, phenyl, substituted phenyl (wherein the substituents are selected from halogen, (C₁-C₆)alkoxy, nitro, NH₂, and


15 (C₁-C₆)alkyl), -(C₁-C₆)alkylphenyl, nitro, -O(CH₂)_qN((C₁-C₆)alkyl)₂ wherein q is 1-3, -NR₁₇R₁₈, 2-thiazole optionally substituted by a (C₁-C₆) alkyl group, -CH=CH-phenyl,

-O-piperazine, pyrrolinyl, and a moiety of the formula:




wherein R₁₇ and R₁₈ are independently selected from H, benzyl, phenyl, (C₁-C₆)alkyl, CH₂CN, CH₂CH₂CN, or R₁₇ and R₁₈ taken together with the nitrogen to which they are attached form a 5-6 membered monocyclic heterocyclic ring optionally containing 1-2 additional nitrogen atoms;

- 5 R₄ is selected from H, (C₁-C₁₀)alkyl (branched or unbranched), (C₁-C₁₀)alkoxy; -O(C₁-C₆)alkylphenyl, CF₃, halo, phenyl, nitro and a moiety of the formula:




or R₂ and R₃ or R₃ and R₄ taken together with the phenyl ring to which they are attached form a moiety selected from those of the formulae:



is CH, O or N and R₂₀ is (C₁-C₃)alkyl or OH;

R₅ is selected from H, amino and amidino;

R₆ is selected from nitro and moieties of the formula:



15


and



wherein:

R₉ and R₁₀ are H or R₉ and R₁₀ taken together form a 5, 6 or 7 membered monocyclic heterocyclic ring optionally containing 1 or 2 additional nitrogens

and optionally substituted wherein the substituent is dimethyl, or R₉ and R₁₀ taken together form a bicyclic heterocyclic ring of the formula;



R₇ is selected from H, amino and nitro;

5 R₈ is selected from H and methyl;

R₁₂ is selected from H and nitro;


R₁₃ is selected from H, nitro, halo, CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and amino;

R₁₄ is selected from H, nitro, halo, CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, amino, amidine, sulfonate, -C(O)O(C₁-C₆)alkyl, and carboxyl;

10 R₁₅ is selected from H, nitro, halo, CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and amino;

R₁₆ is selected from H and nitro;

or R₁₃ and R₁₄ or R₁₄ and R₁₅ taken together form a fused 5 membered heterocyclic ring of the formula:



15 which may be optionally substituted wherein the substituent is selected from amino and methyl;

n is an integer from 0-2;


and the pharmaceutically acceptable salts, esters and prodrug forms thereof.

20 3. A compound having antimicrobial activity according to claim 2 wherein:

R₁-R₄ are as described in claim 2;


R₅ is H;

R₆ is a moiety of the formula:




wherein:

- R₉ and R₁₀ are H or R₉ and R₁₀ taken together form a 5, 6 or 7 membered monocyclic heterocyclic ring optionally containing 1 or 2 additional nitrogens and optionally substituted wherein the substituent is dimethyl, or R₉ and R₁₀ taken together form a bicyclic heterocyclic ring of the formula;



R₇ and R₈ are hydrogen and R₁₂ to R₁₆ are as described above in claim 2; and the pharmaceutically acceptable salts, esters and prodrug forms thereof.

4. A compound having antimicrobial activity selected from those of the formula :




wherein:


R₁ is selected from H, OH, and (C₁-C₆)alkoxy;

R₂ is selected from H, (C₁-C₁₀)alkyl (branched or unbranched), (C₁-C₆)alkoxy; -

O(C₁-C₁₀)alkylphenyl, CF₃, halo, phenyl, nitro and a moiety of the formula:



- R₃ is selected from H, OH, (C₁-C₁₀)alkyl (branched or unbranched), (C₁-C₁₀)alkoxy,
 -O(C₁-C₆)alkylphenyl, CF₃, halo, phenyl, substituted phenyl (wherein the
 substituents are selected from halogen, (C₁-C₆)alkoxy, nitro, NH₂, and (C₁-C₆)alkyl),
 5 - (C₁-C₆)alkylphenyl and and a moiety of the formula:



- R₄ is selected from H, (C₁-C₁₀)alkyl (branched or unbranched), and -OCH₂phenyl;
 R₅ is H;
 R₆ is amidine;
 10 R₇ is H;
 R₈ is H;
 n is 0;
 R₁₂ is H;
 R₁₃ is selected from H, nitro, halo, CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and amino;
 15 R₁₄ is selected from H, nitro, halo, CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy and amino;
 R₁₅ is selected from H and (C₁-C₆) alkoxy, and amino;
 R₁₆ is H;
 and the pharmaceutically acceptable salts, esters and prodrug forms thereof.

- 20 5. The compound of claim 1 wherein R₁ - R₂ are hydrogen, R₃ is phenyl, R₄ - R₅ are hydrogen, R₆ is amidine, and R₇, and R₈ are all hydrogen.

6. The compound of claim 1 wherein R₁ - R₂ are hydrogen, R₃ is -N(phenyl)₂, R₄ - R₅ are hydrogen, R₆ is amidine, and R₇ and R₈ are all hydrogen.


7. The compound of claim 1 wherein $R_1 - R_2$ are hydrogen, R_3 is $-OCH_2$ phenyl, $R_4 - R_5$ are hydrogen, R_6 is amidine, and R_7 and R_8 are all hydrogen.

5 8. The compound of claim 1 wherein R_1 is hydroxy, R_2 is t-butyl, R_3 is hydrogen, R_4 is t-butyl, R_5 is hydrogen, R_6 is amidine, and R_7 and R_8 are all hydrogen.

9. The compound of claim 1 wherein R_1 is hydroxy, R_2 is t-butyl, R_3 is hydrogen, R_4 is hydrogen, R_5 is hydrogen, R_6 is amidine, and R_7 and R_8 are all hydrogen.

10


10. The compound of claim 1 wherein $R_1 - R_2$ are hydrogen; R_3 is a moiety of the formula:



is amidine and R_{15} and R_{16} are hydrogen;

15 $R_4 - R_5$ are hydrogen; R_6 is amidine; and R_7 and R_8 are all hydrogen.


11. The compound of claim 1 wherein $R_1 - R_2$ are hydrogen; R_3 is a moiety of the formula:



20 is hydrogen and R_{15} and R_{16} are hydrogen;


$R_4 - R_5$ are hydrogen; R_6 is amidine; and R_7 and R_8 are all hydrogen.

12. The compound of claim 1 wherein $R_1 - R_2$ are hydrogen; R_3 is a moiety of the formula:



5 $R_4 - R_5$ are hydrogen; R_6 is amidine; and R_7 and R_8 are all hydrogen.


13. The compound of claim 1 wherein $R_1 - R_2$ are hydrogen; R_3 is a moiety of the formula:



10 is chloro and R_{15} and R_{16} are hydrogen;

$R_4 - R_5$ are hydrogen; R_6 is amidine; and R_7 and R_8 are all hydrogen.

14. The compound of claim 1 wherein R_1 is hydrogen; R_2 is a moiety of the formula:



15 and R_{14} , R_{15} and R_{16} are hydrogen;


$R_4 - R_5$ are hydrogen; R_6 is amidine; and R_7 and R_8 are all hydrogen.

15. A pharmaceutical composition for treating bacterial infections comprising an effective amount of a compound selected from claim 1 in association with a
20 pharmaceutically acceptable carrier.

16. A method of treating bacterial infections in mammals by administering to a mammal suffering from such infection a therapeutically effective amount of a compound selected from claim 1.

5

17. A method of preparing a compound of the formula:




Formula I

10 wherein:

R₁ is selected from H, OH, (C₁-C₆)alkoxy;

R₂, R₃ and R₄ are independently selected from H, OH, (C₁-C₁₀)alkyl (branched or unbranched), (C₁-C₁₀)alkoxy, -O(C₁-C₆)alkylphenyl, -OCH₂(4-NO₂phenyl), CF₃, halo, phenyl, substituted phenyl (wherein the substituents are selected from halogen,

15 (C₁-C₆)alkoxy, nitro, NH₂, and (C₁-C₆)alkyl), -(C₁-C₆)alkylphenyl, nitro, -O(CH₂)_qN((C₁-C₆)alkyl)₂ wherein q is 1-3, -NR₁₇R₁₈, 2-thiazole optionally substituted by a (C₁-C₆) alkyl group, -CH=CH-phenyl, -O-piperazine, pyrrolinyl, and a moiety of the formula:




20 wherein n is an integer from 0-2;

R₁₂, R₁₃, R₁₄, R₁₅ and R₁₆ are independently selected from H, nitro, halo, CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and amino, amidine, sulfonate,

-C(O)O(C₁-C₆)alkyl, and carboxyl;

or R₁₃ and R₁₄ or R₁₄ and R₁₅ taken together form a fused 5 membered heterocyclic ring of the formula:




5 which may be optionally substituted wherein the substituent is selected from amino and methyl;

R₁₇ and R₁₈ are independently selected from H, benzyl, phenyl, (C₁-C₆)alkyl, CH₂CN, CH₂CH₂CN, or R₁₇ and R₁₈ taken together with the nitrogen to which they are attached form a 5-6 membered monocyclic heterocyclic ring

10 optionally containing 1-2 additional nitrogen atoms;


or R₂ and R₃ or R₃ and R₄ taken together with the phenyl ring to which they are attached form a moiety selected from those of the formulae:



is CH, O or N and R₂₀ is (C₁-C₃)alkyl or OH;

15 R₅ is selected from H, amino and amidino;

R₆ is selected from nitro and moieties of the formula:




and



20 wherein:

R₉ is hydrogen and R₁₀ is selected from H or OH or R₉ and R₁₀ taken together form a 5, 6 or 7 membered monocyclic heterocyclic ring optionally

containing 1 or 2 additional nitrogens and optionally substituted wherein the substituent is dimethyl, or R₉ and R₁₀ taken together form a bicyclic heterocyclic ring of the formula;



5 R₇ is selected from H, amino and nitro;


R₈ is selected from H and methyl;

and the pharmaceutically acceptable salts, esters and prodrug forms thereof;


which comprises:

reacting a compound of the formula:

10




with a compound of the formula:



in a suitable solvent.

15 18. A method of preparing a compound of the formula:




Formula I

wherein:


20 R₁ is selected from H, OH, (C₁-C₆)alkoxy, and -O(C₁-C₃)COOH;

- R₂, R₃ and R₄ are independently selected from H, OH, (C₁-C₁₀)alkyl (branched or unbranched), (C₁-C₁₀)alkoxy, -O(C₁-C₆)alkylphenyl, -OCH₂(4-NO₂phenyl), CF₃, halo, phenyl, substituted phenyl (wherein the substituents are selected from halogen, (C₁-C₆)alkoxy, nitro, NH₂, and (C₁-C₆)alkyl), -(C₁-C₆)alkylphenyl, nitro, -
- 5 O(CH₂)_qN((C₁-C₆)alkyl)₂ wherein q is 1-3, -NR₁₇R₁₈, 2-thiazole optionally substituted by a (C₁-C₆) alkyl group, -CH=CH-phenyl, -O-piperazine, pyrrolinyl, and a moiety of the formula:




wherein n is an integer from 0-2;

- 10 R₁₂, R₁₃, R₁₄, R₁₅ and R₁₆ are independently selected from H, nitro, halo, CF₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and amino, amidine, sulfonate, -C(O)O(C₁-C₆)alkyl, and carboxyl;
- or R₁₃ and R₁₄ or R₁₄ and R₁₅ taken together form a fused 5 membered heterocyclic ring of the formula:



- 15 which may be optionally substituted wherein the substituent is selected from amino and methyl;
- R₁₇ and R₁₈ are independently selected from H, benzyl, phenyl, (C₁-C₆)alkyl, CH₂CN, CH₂CH₂CN, or R₁₇ and R₁₈ taken together with the nitrogen to
- 20 which they are attached form a 5-6 membered monocyclic heterocyclic ring optionally containing 1-2 additional nitrogen atoms;
- or R₂ and R₃ or R₃ and R₄ taken together with the phenyl ring to which they are attached form a moiety selected from those of the formulae:



is CH, O or N and R₂₀ is (C₁-C₃)alkyl or OH;


R₅ is selected from H, amino and amidino;

R₆ is selected from nitro and moieties of the formula:



wherein:

10 R₉ and R₁₀ are H or R₉ and R₁₀ taken together form a 5, 6 or 7 membered monocyclic heterocyclic ring optionally containing 1 or 2 additional nitrogens and optionally substituted wherein the substituent is dimethyl, or R₉ and R₁₀ taken together form a bicyclic heterocyclic ring of the formula;




R₇ is selected from H, amino and nitro;


15 R₈ is selected from H and methyl;

and the pharmaceutically acceptable salts, esters and prodrug forms thereof; which comprises:

reacting a compound of the formula:



20 with a compound of the formula:



in a suitable solvent.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/18586

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D235/18 A01N43/52 C07D417/10 A01N43/78 C07D405/04
C07D403/04 C07D405/14 A01N43/54 C07D455/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 2 125 790 A (ICI PLC) 14 March 1984 see claim 1	1
X	DE 30 37 188 A (DYNAMIT NOBEL AG) 29 April 1982 see example 3	1
X	EP 0 148 431 A (THOMAE GMBH DR K) 17 July 1985 see claim 1	1
X	WO 96 06831 A (CANCER SOC AUCKLAND DIV NZ INC ;CIRCADIAN PHARM AU PTY LTD (AU); D) 7 March 1996 see claim 1	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

26 November 1998

Date of mailing of the international search report

14/12/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gettins, M

INTERNATIONAL SEARCH REPORT

Int. J. Application No.

PCT/US 98/18586

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 353 606 A (KONISHIROKU PHOTO IND) 7 February 1990 see example 4 ---	1
X	US 4 982 112 A (OKAZAKI MASAKI ET AL) 1 January 1991 see column 6; example 8 ---	1
X	EP 0 511 187 A (UCB SA) 28 October 1992 see claim 1; examples 10,15 ---	1
X	ERTEPINAR ET AL: "A QSAR study of the biological activities of some benzimidazoles and imidazopyridine against Bacillus subtilis" EUR.J.MED.CHEM., vol. 30, 1995, pages 171-175, XP002085872 see examples 2,5; table I ---	1-18
X	COBURN ET AL: "Substituted 2-(2-Hydroxyphenyl)benzimidazoles as potential agents for the control of periodontal disorders." J.MED.CHEM., vol. 87, no. 30, 1987, pages 205-208, XP002085873 see example 9D; table I ---	1-18
X	JOSHI ET AL: "Synthesis of some novel 1,3,4-oxadiazoles and 5-oxo-imidazolines as potent biologically active agents" HETEROCYCL.COMMUN., vol. 3, no. 2, 1997, pages 169-174, XP002085874 see example 2 -----	1-18

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int'l Application No

PCT/US 98/18586

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2125790 A	14-03-1984	NONE	
DE 3037188 A	29-04-1982	NONE	
EP 0148431 A	17-07-1985	DE 3346575 A	04-07-1985
		AU 3704284 A	04-07-1985
		DD 232697 A	05-02-1986
		DK 597484 A	24-06-1985
		FI 845084 A	24-06-1985
		JP 60156676 A	16-08-1985
		PT 79718 B	30-12-1986
WO 9606831 A	07-03-1996	AU 695936 B	27-08-1998
		AU 3247595 A	22-03-1996
		EP 0777656 A	11-06-1997
EP 0353606 A	07-02-1990	JP 2040634 A	09-02-1990
		JP 2700475 B	21-01-1998
		US 5112532 A	12-05-1992
US 4982112 A	01-01-1991	JP 1944545 C	23-06-1995
		JP 6079125 B	05-10-1994
		JP 62210430 A	16-09-1987
		JP 1985250 C	25-10-1995
		JP 7011646 B	08-02-1995
		JP 62210432 A	16-09-1987
		DE 3707835 A	17-09-1987
		US 5115337 A	19-05-1992
EP 0511187 A	28-10-1992	AT 145398 T	15-12-1996
		DE 69215266 D	02-01-1997
		DE 69215266 T	28-05-1997
		DK 511187 T	07-04-1997
		ES 2096740 T	16-03-1997
		GR 3022603 T	31-05-1997